



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

HONDURAS.

*Sanitary report of Puerto Cortez and Ceiba.*PORT TAMPA, FLA., *February 23, 1898.*

SIR: I have the honor to report as per your request, the sanitary condition of Honduras. Unfortunately I missed seeing Dr. Austen at Puerto Cortez, and the information I gained there was from observation and what I could collect from the residents. This place has a population, with the adjacent settlement, of about 1,200, and is fairly healthy. Malarial diseases are very prevalent there, and although not very malignant, are quite serious, and quinine is taken freely. Ceiba, a coast town about 60 miles south of Puerto Cortez, I found very unhealthy; the principal diseases are malarial and intestinal. The sanitary condition there is very bad; the water used is surface. Intestinal diseases are very malignant and the mortality very high, especially among children.

Very respectfully, yours,

G. H. ALTREE,

Acting Assistant Surgeon, U. S. M. H. S.

STATISTICAL REPORTS.

BAHAMAS—*Dunmore Town*.—Two weeks ended February 25, 1898. Estimated population, 1,472. One death.

Governors Harbor.—Two weeks ended February 26, 1898. Estimated population, 1,500. One death. No deaths from contagious disease.

Green Turtle Cay—Abaco.—Two weeks ended February 24, 1898. Estimated population, 3,900. No deaths from contagious disease.

CANADA—*Nova Scotia—Windsor*.—Month of February, 1898. Estimated population, 3,000. Total deaths, 2, including 1 from phthisis pulmonalis.

CHILE—*Antofagasta*.—Estimated population, 14,000. Total deaths, 47, including 3 from phthisis pulmonalis.

FRANCE—*Roubaix*.—Month of January, 1898. Estimated population, 124,447. Total deaths, 281, including typhus fever, 1; enteric fever, 3; diphtheria, 4; and measles, 45.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended February 19 correspond to an annual rate of 20.4 a thousand of the aggregate population, which is estimated at 11,218,378. The highest rate was recorded in Wolverhampton, viz, 29.0, and the lowest in Preston, viz, 13.9.

London.—One thousand eight hundred and fifty-six deaths were registered during the week, including measles, 100; scarlet fever, 9; diphtheria, 33; whooping cough, 53; enteric fever, 8, and diarrhea and dysentery, 8. The deaths from all causes correspond to an annual rate of 21.5 a thousand. In greater London 2,392 deaths were registered, corresponding to an annual rate of 19.4 a thousand of the population. In the "outer ring" the deaths included 18 from diphtheria, 13 from measles, 4 from scarlet fever, and 9 from whooping cough.

Ireland.—The average annual death rate represented by the deaths